arXiv:1512.00097v6 [math.AG] 8 Apr 2016

BASIC FINITE ÉTALE EQUIVALENCE RELATIONS

YING ZONG

1. *Introduction.*

Let S be an algebraic space, A an S-abelian algebraic space, X an A-torsor on S for the étale topology and L a finite étale S-equivalence relation on X. Thus, X is representable by an S-algebraic space and its quotient, in the sense of étale topology, by the finite étale S-equivalence relation L, is representable by a proper and smooth S-algebraic space Y with geometrically irreducible fibers. In the following we define the property for L to be a *basic* finite étale equivalence relation and show that this property is remarkably rigid and our first goal is to prove :

Theorem 1.1. — *Let $f : Y \to S$ be a proper smooth morphism of algebraic spaces with S connected. Assume that for one geometric point t of S there is a finite étale t-morphism from a t-abelian variety A'_t onto $Y_t = f^{-1}(t)$ with degree $[A'_t : Y_t]$ prime to the residue characteristics of S.*

Then :

a) There exist an S-abelian algebraic space A, an A-torsor X on S for the étale topology and a basic finite étale S-equivalence relation L on X such that Y is S-isomorphic to the quotient X/L.

b) Let (A, X, L) be as in a). Then X is a f^*G-torsor on Y for the étale topology where the S-group $G = f_*\underline{\mathrm{Aut}}_Y(X)$ is a finite étale S-algebraic space. The functor

$$(A', X', L') \mapsto P' = f_*\underline{\mathrm{Isom}}_Y(X, X')$$

induces a bijection from the set of all triples as in a) up to isomorphisms onto $H^1(S, G)$.

The proof is in §9.

When f degenerates to $\overline{f} : \overline{Y} \to \overline{S}$ where $\overline{f}$ is separated of finite type and universally open with $\overline{S}$ locally noetherian, then a geometric fiber $\overline{f}^{-1}(\overline{s})$, if it is non-empty and does not have uniruled irreducible components, is irreducible; see §10, which, as a continuation of [4] and our main objective, contains further support for the *Principle* : *Non-uniruled abelian degenerations are almost non-degenerate.*

2. *Definition.*

Definition 2.1. — *Let S be an algebraic space, A an S-abelian algebraic space, X an A-torsor on S for the étale topology and L a finite étale S-equivalence relation on X. We say that L is basic at a geometric point s of S provided the following condition holds* :

If E is a finite étale s-subgroup of A_s such that the morphism

$$E \times_s X_s \to X_s \times_s X_s,\ (e, x) \mapsto (e + x, x)$$

factors through the graph of L_s, then $E = 0$.

We say that L is basic if it is basic at every geometric point of S.

Thus, put in words, L is basic at s if and only if no sub-equivalence relation of $L(s)$ other than the identity is generated by a translation. Let Y denote the quotient X/L. The graph of L is then identical to $X \times_Y X$. With the above notations, saying that the morphism

$$E \times_s X_s \to X_s \times_s X_s,\ (e, x) \mapsto (e + x, x)$$

factors through the graph of L_s amounts to saying that the translations of X_s defined by the elements of $E(s)$ are Y_s-automorphisms.

2.2. Let $s' \to s$ be a morphism of geometric points of S. The base change

$$L_s \mapsto L_s \times_s s'$$

establishes a bijection between the collection of finite étale s-equivalence relations on X_s which are basic at s and the collection of finite étale s'-equivalence relations on $X_{s'}$ which are basic at s'. Indeed, for every proper s-algebraic space Z, the functor of base change by $s' \to s$ is an equivalence from the category of finite étale Z-algebraic spaces onto the category of finite étale $Z_{s'}$-algebraic spaces (SGA 4 XII 5.4). This same base change provides also an equivalence between the category of finite étale s-subgroups of A_s and the category of finite étale s'-subgroups of $A_{s'}$. In particular, a given finite étale S-equivalence relation L on X is basic at s if and only if it is basic at s'. The collection of basic finite étale equivalence relations on X_s and on $X_{s'}$ are in canonical bijective correspondence. One deduces also that if L is basic, so is $L_{S'}$ for every base change $S' \to S$. When $S' \to S$ is surjective, the condition that L be basic is then equivalent to that $L_{S'}$ be basic.

3. *Automorphisms I.*

We study now the Y-automorphisms of X.

Lemma 3.1. — *Keep the notations of* (2.1). *With each S-morphism $q : X \to X$ there is associated a unique S-group homomorphism $p : A \to A$ such that q is p-equivariant.*

Proof. One may assume the torsor X to be trivial, as the existence of a unique homomorphism p is an étale local question on S.

Each S-morphism $q : A \to A$ is the unique composite of a translation $(x \mapsto x+q(0))$ and an S-group homomorphism $p : A \to A$ ("Geometric Invariant Theory" 6.4). It is clear that q is p-equivariant : for every two local S-sections a, x of A, one has

$$q(a + x) = p(a) + q(x).$$

□

Lemma 3.2. — *Keep the notations of* (2.1). *Assume that L is basic. Then every Y-endomorphism of X is a Y-automorphism. If X admits furthermore an S-section x, the only Y-automorphism of X preserving x is the identity morphism.*

Proof. Every Y-morphism $X \to X$ is finite étale, as X is finite étale over Y. Let q be such a Y-endomorphism of X. By (3.1), q is equivariant relative to a unique S-group homomorphism $p : A \to A$. And, p is finite étale, since it is étale locally on S isomorphic to q. In particular, $E = \mathrm{Ker}(p)$ is a finite étale S-subgroup of A. Observe that the morphism

$$E \times_S X \to X \times_S X,\ (e, x) \mapsto (e + x, x)$$

factors through the graph of L. That is to say, for every local S-section e (resp. x) of E (resp. X), $(e + x, x)$ is a local S-section of $X \times_Y X$. Indeed, as q is a Y-endomorphism of X, the sections x and $q(x)$ (resp. $e + x$ and $q(e + x) = p(e) + q(x) = q(x)$) have the same image in Y.

So $E = 0$, as L is basic. So p, hence q as well, is an isomorphism.

Let $Z = \mathrm{Ker}(q, \mathrm{Id}_X)$, which is an open and closed sub-Y-algebraic space of X. If $q(x) = x$, that is, if $x \in Z(S)$, then $Z = X$ and $q = \mathrm{Id}_X$, since X has geometrically irreducible S-fibers.

□

Lemma 3.3. — *For $i = 1, 2$, let $X_i, L_i, Y_i = X_i/L_i$ be as in* (2.1), *with L_i basic, let $x_i \in X_i(S)$ and let $y_i \in Y_i(S)$ be the image of x_i. Let $r : Y_1 \to Y_2$ be an S-isomorphism satisfying $r(y_1) = y_2$.*

Then there exists a unique r-isomorphism $h : X_1 \to X_2$ satisfying $h(x_1) = x_2$.

Proof. Identify Y_1 with Y_2 by r. Write $Y = Y_1 = Y_2$ and $y = y_1 = y_2$. The uniqueness of h follows from (3.2). Thus, the existence of h is an étale local question on S.

One may assume that S is strictly local with closed point s. For, if the lemma is proven under this assumption, then h exists in an étale

neighborhood of s in S by a "passage à la limite projective" (EGA IV 8). As the closed immersion $Y_s \hookrightarrow Y$ induces an equivalence between the category of finite étale Y-algebraic spaces and the category of finite étale Y_s-algebraic spaces (SGA 4 XII 5.9 bis), one may further assume that $S = s$ is the spectrum of a separably closed field k.

Let X be an algebraic space which is connected and finite étale Galois over Y and which dominates X_1 and X_2, say by $q_i : X \to X_i$, $i = 1, 2$. By SGA 1 XI 2, X is a (trivial) torsor under a k-abelian variety A and q_i is equivariant relative to a unique étale k-group homomorphism $p_i : A \to A_i$, $i = 1, 2$.

With each element $g \in \mathrm{Gal}(X/Y) = G$, there is associated a unique group automorphism $a(g)$ of the k-abelian variety A such that g is $a(g)$-equivariant. The map $g \mapsto a(g)$ is a homomorphism of groups and its kernel $a^{-1}(1) = E$ consists of translations by elements of $A(k)$.

By definition, $\mathrm{Ker}(p_1)$ and $\mathrm{Ker}(p_2)$ are k-subgroups of E_k. Hence, X/E is dominated by X_1 and by X_2. So $E_k = \mathrm{Ker}(p_1) = \mathrm{Ker}(p_2)$, as L_1 and L_2 are basic.

There exist therefore a unique Y-isomorphism $q : X_1 \to X_2$ and a unique isomorphism of k-abelian varieties $p : A_1 \to A_2$ such that q is p-equivariant and such that $qq_1 = q_2$, $pp_1 = p_2$.

Identify X_1 with X_2 by q and identify A_1 with A_2 by p. Then, $X_1 = X_2 = X/E$ is Galois over Y with Galois group G/E. Clearly, a unique element h of G/E transforms x_1 to x_2.

□

Proposition 3.4. — *Keep the notations of* (2.1)*. Assume that L is basic. Let $f : Y \to S$ denote the structural morphism.*

Then the S-group $G = f_*\underline{\mathrm{Aut}}_Y(X)$ is a finite étale S-algebraic space and the canonical morphism

$$u : G \times_S X \to X \times_Y X,\ (g, x) \mapsto (g(x), x)$$

is an isomorphism.

Proof. The finite étale Y-group $N = \underline{\mathrm{Aut}}_Y(X)$, considered as a proper smooth S-algebraic space, has Stein factorization $N \to f_*N \to S$ (SGA 1 X 1.2). By *loc.cit.*, $f_*N \to S$ is finite étale and the formation of the Stein factorization commutes with every base change $S' \to S$. It remains only to verify that $u \times_S s$ is an isomorphism for each geometric point s of S. One can thus assume that S is the spectrum of a separably closed field. Then, by the proof of (3.3), $X \to Y$ is Galois and hence u is an isomorphism.

□

Proposition 3.5. — *For $i = 1, 2$, let $X_i, L_i, Y_i = X_i/L_i$ be as in* (2.1) *with L_i basic, let $f_i : Y_i \to S$ be the structural morphism, let $G_i = f_{i*}\underline{\mathrm{Aut}}_{Y_i}(X_i)$ and let $r : Y_1 \to Y_2$ be an S-isomorphism. Write X_i', L_i', Y_i', r' for the base change of X_i, L_i, Y_i, r by a morphism $S' \to S$. Then the sheaf on* (Sch/S) *for the étale topology, $S' \mapsto \mathrm{Isom}_{r'}(X_1', X_2')$, is a (G_2, G_1)-bitorsor. The map*

$$\mathrm{Isom}_r(X_1, X_2) \to \mathrm{Isom}_{r'}(X_1', X_2'),\ h \mapsto h \times_S S'$$

is a bijection if $S' \to S$ is 0*-acyclic* (SGA 4 XV 1.3).

Proof. Identify Y_1 with Y_2 by r, write $Y = Y_1 = Y_2$ and $f = f_1 = f_2$. Let $I = \underline{\mathrm{Isom}}_Y(X_1, X_2)$, which is a finite étale Y-algebraic space. By SGA 1 X 1.2, f_*I is a finite étale S-algebraic space and its formation commutes with every base change $S' \to S$ and it is a (G_2, G_1)-bitorsor by the same argument of (3.4). When $u : S' \to S$ is 0-acyclic, the adjunction morphism

$$f_*I \to u_*u^*f_*I$$

is an isomorphism and in particular

$$\Gamma(Y, I) = \Gamma(S, f_*I) \to \Gamma(Y', I') = \Gamma(S, u_*u^*f_*I)$$

is a bijection.

□

4. *Factorization over a separably closed field.*

Proposition 4.1. — *Over a separably closed field k, every finite étale surjective k-morphism from a k-abelian variety A' to a k-algebraic space Y, $A' \to Y$, factors up to k-isomorphisms in a unique way as the composite of an étale isogeny of k-abelian varieties $A' \to A$ and a quotient $A \to A/L = Y$ by a basic finite étale k-equivalence relation L on A.*

Proof. Let $A' \to Y$ be dominated by a finite étale Galois k-morphism $A'' \to Y$ with A'' connected. By SGA 1 XI 2, A'' may be endowed with a k-abelian variety structure so that the projection $A'' \to A'$ is an étale k-group homomorphism. Let E'' denote the kernel of this homomorphism.

Let E consist of the elements of $G = \mathrm{Gal}(A''/Y)$ which are translations of A'', namely, of the form $x \mapsto x + a$, with $a \in A''(k)$. Evidently, E is a subgroup of G, $E \supset E''(k)$ and the quotient $A = A''/E$ inherits a k-abelian variety structure from that of A''.

The factorization

$$A' = A''/E'' \to A = A''/E \to Y = A''/G$$

is then as desired. One argues as in (3.3) that such a factorization is unique up to k-isomorphisms.

□

5. *Basicness is an open and closed property.*

Proposition 5.1. — *Keep the notations of* (2.1). *Let*

$$F : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$$

be the following sub-functor of the final functor :

For an S-scheme S', $F(S') = \{\emptyset\}$, if $L \times_S S'$ is basic, and $F(S') = \emptyset$, otherwise.

Then F is representable by an open and closed sub-algebraic space of S.

Lemma 5.2. — *Keep the notations of* (2.1). *Assume that S is the spectra of a discrete valuation ring with generic point t and closed point s.*

Then the following conditions are equivalent :

1) *L_s is basic.*

2) *L_t is basic.*

Proof. One may assume that S is strictly henselian.

Assume 1). The finite étale morphism $X_s \to X_s/L_s$ is Galois (3.4). So by SGA 4 XII 5.9 bis, X is Galois over $X/L = Y$, as S is strictly henselian and $Y \to S$ is proper. Let $G = \mathrm{Gal}(X/Y)$. Each element $g \in G$ is equivariant relative to a unique group automorphism $a(g)$ of the S-abelian algebraic space A. The homomorphism $g \mapsto a(g)_s$ is injective, as L_s is basic. It follows that $g \mapsto a(g)_t$ is injective. For, the specialization homomorphism

$$\mathrm{Aut}_t(A_t) \xleftarrow{\sim} \mathrm{Aut}_S(A) \to \mathrm{Aut}_s(A_s)$$

is injective. So L_t is basic.

Assume 2). Let $\overline{t}$ be the spectrum of a separable closure of $k(t)$. By (3.4), $X_{\overline{t}} \to Y_{\overline{t}}$ is Galois with Galois group, say G. Replacing if necessary S by its normalization S' in a finite sub-extension of $k(\overline{t})/k(t)$, X by $X \times_S S'$, Y by $Y \times_S S'$ and L by $L \times_S S'$, one may assume that X_t is a G-torsor on Y_t and that L_t is defined by the G-action on X_t. Now X being the S-Néron model of X_t, each t-automorphism of X_t uniquely extends to an S-automorphism of X. In particular, the G-action on X_t extends to an action on X with graph evidently equal to that of L. For each $g \in G$, let $a(g)$ be the unique S-automorphism of the S-abelian algebraic space A relative to which $g : X \to X$ is equivariant. The

homomorphism $g \mapsto a(g)_t$ is injective, as L_t is basic. Then $g \mapsto a(g)_s$ is injective, for the specialization homomorphism

$$\mathrm{Aut}_t(A_t) \widetilde{\leftarrow} \mathrm{Aut}_S(A) \to \mathrm{Aut}_s(A_s)$$

is injective. So L_s is basic.

□

Lemma 5.3. — *Keep the notations of* (2.1). *Assume that L is basic. Assume that S is an affine scheme and that (S_i) is a projective system of affine noetherian schemes, indexed by a co-directed set I, with limit S.*

Then there exist an index $i \in I$, an S_i-abelian algebraic space A_i, an A_i-torsor X_i on S_i for the étale topology and a basic finite étale S_i-equivalence relation L_i on X_i such that $A = A_i \times_{S_i} S$, $X = X_i \times_{S_i} S$ and $L = L_i \times_{S_i} S$.

Proof. By the technique of "passage à la limite projective" (EGA IV 8, 9), there exist $i_o \in I$ and $A_{i_o}, X_{i_o}, L_{i_o}$ as desired except possibly the property of being *basic*. By (5.2), for each $j \geq i_o$, $L_{i_o} \times_{S_{i_o}} S_j$ is basic at precisely the points of an open and closed sub-scheme S'_j of S_j. The projection $S \to S_j$ factors through S'_j, as L is basic. Now $(S_j - S'_j)_{j \geq i_o}$ is a projective system of affine noetherian schemes with empty limit. So $S_i = S'_i$ for some $i \geq i_o$.

□

5.4. *Proof of* (5.1).

The question being an étale local question on S, one may assume that S is a scheme, then affine and by (5.3) noetherian. The claim is now immediate by (5.2).

□

6. *Automorphisms II.*

We study the S-automorphisms of Y.

Lemma 6.1. — *Keep the notations of* (2.1). *Assume that L is basic. Let $p : X \to X/L = Y$ denote the projection. Let r be an S-automorphism of Y.*

Then there is at most one S-section a of A which verifies $rp = pT_a$, where $T_a : x \mapsto a + x$ is the S-automorphism of X, the translation by a.

Proof. If $rp = pT_a = pT_b$ holds for two S-sections a, b of A, then T_{a-b} acts on X as a Y-automorphism. In particular, T_{a-b} is locally on S of finite order. So $T_{a-b} = \mathrm{Id}_X$ and $a = b$, as L is basic.

□

Lemma 6.2. — *Keep the notations and assumptions of* (6.1).

Let $F : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$ *be the following sub-functor of the final functor* :

For an S*-scheme* S', $F(S') = \{\emptyset\}$, *if* $r_{S'}p_{S'} = p_{S'}T_{a'}$ *for an* S'*-section* a' *of* $A_{S'}$, *and* $F(S') = \emptyset$, *otherwise.*

Then F *is representable by an open and closed sub-algebraic space of* S.

Proof. The question being by (6.1) an étale local question on S, one may assume S to be a scheme, then affine and by (5.3) noetherian.

— *The functor* F *verifies the valuative criterion of properness* :

Namely, given an S-scheme S' which is the spectra of a discrete valuation ring and which has generic point t', then $F(S') = F(t')$. For, if $r_{t'}p_{t'} = p_{t'}T_{a_{t'}}$ holds for some point $a_{t'} \in A(t')$, then $a_{t'}$ extends uniquely to an S'-section a' of $A_{S'}$ and the equation $r_{S'}p_{S'} = p_{S'}T_{a'}$ holds.

— *The functor* F *is formally étale* :

Namely, $F(S') = F(S'')$ for every nilpotent S-immersion $S'' \hookrightarrow S'$. Indeed, assume $r_{S''}p_{S''} = p_{S''}T_{a''}$ holds for some section $a'' \in A(S'')$. As both p and rp are étale, there is a unique S'-automorphism T' of $X_{S'}$ such that T' restricts to $T_{a''}$ on $X_{S''}$ and such that $r_{S'}p_{S'} = p_{S'}T'$ holds. This T' is equivariant with respect to a unique S'-group automorphism φ' of $A_{S'}$. As the S-group $\underline{\mathrm{Aut}}_S(A)$ is unramified over S and as φ' restricts to the identity automorphism of $A_{S''}$, it follows that $\varphi' = \mathrm{Id}_{A_{S'}}$. So T' is of the form $T_{a'}$ for an S'-section $a' \in A(S')$.

It is now evident that $F \to S$ is representable by an open and closed immersion.

□

Proposition 6.3. — *Keep the notations of* (2.1). *Assume that* L *is basic. Let* $p : X \to X/L = Y$ *denote the projection and* $f : Y \to S$ *the structural morphism.*

Then :

a) The S*-subgroup* R *of* $\underline{\mathrm{Aut}}_S(Y)$ *defined to consist of the local* S*-automorphisms* r *satisfying* $rp = pT_a$ *for some local* S*-sections* a *of* A *is open and closed in* $\underline{\mathrm{Aut}}_S(Y)$.

b) The homomorphism $r \mapsto a$ *is a closed immersion of* R *into* A. *In particular,* R *is commutative and representable by a proper* S*-algebraic space.*

c) An S-section a of A lies in the image of R if and only if $T_a g = g T_a$ for all local S-sections g of $G = f_* \underline{\mathrm{Aut}}_Y(X)$.

Proof. Note that $R \to A$, $r \mapsto a$, is a well-defined morphism by (6.1). This morphism is a monomorphism because p, being étale surjective, is an epimorphism in the category of S-algebraic spaces.

The claim that the sub-S-group functor $R \subset \underline{\mathrm{Aut}}_S(Y)$ is open and closed is a rephrase of (6.2), hence a).

If an S-section a of A satisfies $T_a g = g T_a$ for every local S-section g of G, then T_a, being G-equivariant, defines by passing to the G-quotient an S-automorphism r of $X/G = Y$ (3.4). One has thus $rp = pT_a$ by construction. Conversely, the identity $rp = pT_a$ implies that, for every local S-section g of G, one has $rp = rpg^{-1} = pgT_a g^{-1}$ and then by (6.1) $gT_a g^{-1} = T_a$. The characterization in c) therefore follows and it implies evidently that the monomorphism $R \hookrightarrow A$, $r \mapsto a$, is a closed immersion, hence b).

□

7. *Rigidity of torsors and albanese.*

Lemma 7.1. — *Let S be the spectra of a discrete valuation ring and t (resp. s) a geometric generic (resp. closed) point of S. Let X be a proper smooth S-algebraic space.*

Then the following conditions are equivalent :

1) *$X \times_S s$ has an s-abelian variety structure.*

2) *$X \times_S t$ has a t-abelian variety structure.*

Proof. One may assume that S is strictly henselian. Then X has S-sections by EGA IV 17.16.3. Fix an S-section o of X. When 1) (resp. 2)) holds, after a translation of the origin, $X \times_S s$ (resp. $X \times_S t$) has an abelian variety structure over s (resp. t) with o_s (resp. o_t) as its zero section. Under either assumption, X has geometrically irreducible S-fibers and admits a unique S-abelian algebraic space structure with o being the zero section ("Geometric Invariant Theory" 6.14).

□

Proposition 7.2. — *Let S be an algebraic space and X a proper smooth S-algebraic space. Let $T : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$ be the following sub-functor of the final functor* :

For an S-scheme S', $T(S') = \{\emptyset\}$, if $X \times_S s'$ has an s'-abelian variety structure for each geometric point s' of S', and $T(S') = \emptyset$, otherwise.

Then T is representable by an open and closed sub-algebraic space of S. And $X \times_S T = X_T$ is a $\mathrm{Pic}^o_{\mathrm{P}_T/T}$-torsor on T for the étale topology where $\mathrm{P}_T = \mathrm{Pic}^o_{X_T/T}$ is a T-abelian algebraic space.

Proof. From (7.1) and by a "passage à la limite", one deduces that $T \to S$ is representable by an open and closed immersion. Replacing S by T, one may assume $T = S$. Let $S' \to S$ be a smooth morphism with geometrically irreducible fibers such that $X' = X \times_S S'$ admits an S'-section e'. One may for instance take $S' = X$ and e' to be the diagonal section $\Delta_{X/S}$. Notice that in $S'' = S' \times_S S'$ the only open and closed neighborhood of the diagonal is S''. By "Geometric Invariant Theory" 6.14, on X' there is a unique S'-abelian algebraic space structure with zero section e'. It follows that $\mathrm{P} = \mathrm{Pic}^\tau_{X/S}$, being representable by an S-algebraic space (Artin), is an S-abelian algebraic space and that $\mathrm{Pic}^\tau_{X/S} = \mathrm{Pic}^o_{X/S}$, as one verifies after the *fppf* base change $S' \to S$ on the S'-abelian algebraic space X'. Let the dual S-abelian algebraic space of P be A. Let p_1, p_2 be the two projections of S'' onto S' and $u : p_1^* X' \to p_2^* X'$ the descent datum on X' relative to $S' \to S$ corresponding to X. By (3.1)+(7.3) u is equivariant with respect to a unique S''-group automorphism a of $A'' = A \times_S S''$. It suffices evidently to show that $a = 1$. Now the S-group $\underline{\mathrm{Aut}}_S(A)$ being unramified and separated over S, the relation "$a = 1$" is an open and closed relation on S'' and holds on the diagonal and so holds.

□

Lemma 7.3. — *Let S be an algebraic space, A an S-abelian algebraic space and X an A-torsor on S for the étale topology.*

Then there exists a canonical isomorphism

$$X \overset{A}{\wedge} \mathrm{Pic}_{A/S} \xrightarrow{\sim} \mathrm{Pic}_{X/S}$$

which induces isomorphisms

$$\mathrm{Pic}^o_{A/S} = X \overset{A}{\wedge} \mathrm{Pic}^o_{A/S} \xrightarrow{\sim} \mathrm{Pic}^o_{X/S},$$

$$\mathrm{NS}_{A/S} = X \overset{A}{\wedge} \mathrm{NS}_{A/S} \xrightarrow{\sim} \mathrm{NS}_{X/S}.$$

Proof. This is "Faisceaux amples sur les schémas en groupes et les espaces homogènes" XIII 1.1, by which it is justified to call A the *albanese* of X.

□

8. *Specialization and descent.*

Proposition 8.1. — *Let S be the spectra of a discrete valuation ring and t (resp. s) a geometric generic (resp. closed) point of S. Let Y be a proper smooth S-algebraic space. Suppose that Y_t is the quotient of a t-abelian variety A_t by a basic finite étale t-equivalence relation such that $[A_t : Y_t]$ is prime to the characteristic of $k(s)$.*

Then Y_s is the quotient of an s-abelian variety by a basic finite étale s-equivalence relation.

Proof. One may assume that S is strictly henselian with closed point s. Recall that by (3.4) A_t is Galois over Y_t. As Y is proper smooth over S, S strictly henselian and $G = \mathrm{Gal}(A_t/Y_t)$ of order prime to the characteristic of $k(s)$, the specialization morphism

$$H^1(Y, G) \to H^1(Y_t, G)$$

is by SGA 4 XVI 2.2 a bijection. One finds thus a G-torsor on Y for the étale topology, $A \to Y$, which specializes to $A_t \to Y_t$ at t. In particular, A is proper smooth over S and has S-sections by EGA IV 17.16.3 and has by (7.2) an S-abelian algebraic space structure. The finite étale S-equivalence relation on A of graph $A \times_Y A$ is basic because it is basic at t (5.1).

□

Proposition 8.2. — *Let S be an algebraic space. Let Y be an algebraic space which is proper flat of finite presentation over S.*

Let $U : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$ be the following sub-functor of the final functor :

For an S-scheme S', $U(S') = \{\emptyset\}$, if $Y \times_S s'$ is the quotient of s'-abelian variety by a finite étale s'-equivalence relation for each geometric point s' of S', and $U(S') = \emptyset$, otherwise.

Then $U \to S$ is representable by an open immersion of finite presentation. There exist a U-abelian algebraic space A, an A-torsor X on U for the étale topology and a basic finite étale U-equivalence relation L on X such that $Y \times_S U$ is U-isomorphic to the quotient X/L.

Proof. Observe that, by (4.1), for a given geometric point s of S, $U(s) = \{\emptyset\}$ if and only if Y_s is the quotient of an s-abelian variety by a *basic* finite étale s-equivalence relation.

— *Reduction to the case where Y is S-smooth* :

Consider the functor $V : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$:

For an S-scheme S', $V(S') = \{\emptyset\}$, if $Y \times_S s'$ is smooth over s' for each geometric point s' of S', and $V(S') = \emptyset$, otherwise.

It is evident that $V \to S$ is representable by an open immersion of finite presentation, that $Y \times_S V$ is proper and smooth over V and that V contains U as a sub-functor. Restricting to V, one may assume that $S = V$, namely, that Y is smooth over S.

— *Assume that Y is S-smooth. Then $U \to S$ is representable by an open immersion of finite presentation. There exist an étale surjective morphism $U' \to U$, a U'-abelian algebraic space A', an A'-torsor X' on U' for the étale topology and a basic finite étale U'-equivalence relation L' on X' such that $Y \times_S U'$ is U'-isomorphic to the quotient X'/L'* :

The question being an étale local question on S, one may assume that S is a scheme, then affine, then by (5.3) noetherian and strictly local with closed point s and finally that $U(s) = \{\emptyset\}$, i.e., that Y_s is the quotient of an s-abelian variety A_s by a basic finite étale s-equivalence relation L_s. By SGA 4 XII 5.9 bis, there is a finite étale S-morphism $A \to Y$ which specializes to $A_s \to A_s/L_s = Y_s$ at s. This algebraic space A is in particular proper smooth over S and has closed fiber the abelian variety A_s. As S is strictly local, A admits S-sections (EGA IV 17.16.3). So by (7.2) A has an S-abelian algebraic space structure. The finite étale S-equivalence relation L on A of graph $A \times_Y A$, basic at s, is basic (5.1).

— *Assume that Y is S-smooth. Then up to an étale localization on U' there exists a descent datum on (A', X', L') relative to $U' \to U$* :

Restricting to the open sub-algebraic space U, one may assume that $U = S$. Write S' for U' and Y' for $Y \times_S S'$. It suffices to prove the existence of a finite étale Y-algebraic space X such that the finite étale Y'-algebraic space $\underline{\mathrm{Isom}}_{Y'}(X \times_Y Y', X')$ is surjective over Y', for then X is an A-torsor on S for the étale topology (7.2) where the S-abelian algebraic space A satisfies $\mathrm{Pic}^o_{A/S} = \mathrm{Pic}^o_{X/S}$ (7.3) and $X \times_Y X$ is the graph of a basic finite étale S-equivalence relation L on X.

i) *Case where Y has an S-section y* :

By an étale localization on S', one may assume that there is an S'-section x' of X' which is mapped to $y' = y \times_S S'$ by $X' \to Y'$. Let $S'' = S' \times_S S'$ and p_1, p_2 the two projections of S'' onto S'. By (3.3), there exists a unique $(Y \times_S S'')$-isomorphism $h : p_1^* X' \to p_2^* X'$ which transforms $p_1^*(x')$ to $p_2^*(x')$. That is, h is a gluing datum on (X', x') relative to $S' \to S$. By (3.3) again, h is a descent datum. This descent provides a finite étale Y-algebraic space X which verifies $X \times_Y Y' = X'$ and which is equipped with an S-section x having image y in $Y(S)$.

ii) *General case* :

By the second projection, $Y \times_S Y = Y_1$ has a Y-algebraic space structure which admits the Y-section $\Delta_{Y/S}$. One finds by i) a finite étale Y_1-algebraic space X_1 such that the finite étale Y_1'-algebraic space

$$\underline{\mathrm{Isom}}_{Y_1'}(X_1 \times_{Y_1} Y_1', X' \times_{Y'} Y_1')$$

is surjective over Y_1', where $Y_1' := Y \times_S Y' = Y_1 \times_Y Y'$. Let the Stein factorization of the proper smooth Y-algebraic space X_1 be

$$X_1 \to X \to Y$$

and let c be the canonical morphism

$$c : X_1 \to X \times_Y Y_1.$$

By SGA 1 X 1.2, the morphism $X \to Y$ is finite étale and the formation of the Stein factorization commutes with every base change $T \to S$. It suffices to show that c is an isomorphism, for then

$$\underline{\mathrm{Isom}}_{Y_1'}(X_1 \times_{Y_1} Y_1', X' \times_{Y'} Y_1') = \underline{\mathrm{Isom}}_{Y'}(X \times_Y Y', X') \times_{Y'} Y_1'$$

and X is the sought after Y-algebraic space. This amounts to showing that $c \times_S s$ is an isomorphism for each geometric point s of S. One may thus assume that S is the spectrum of a separably closed field. Then S', being étale surjective over S, is a non-empty disjoint union of S. Index these components of S' as S_i, $i \in \pi_o(S') = \Pi$, write $X' = \sum_{i \in \Pi} X_i$, fix a point $y \in Y(S)$ and choose a point $x_i \in X_i(S_i)$ above y for each $i \in \Pi$. These (X_i, x_i)'s are all mutually Y-isomorphic by (3.3). Clearly, c is an isomorphism.

□

9. *Proof of Theorem* 1.1.

a) Let the open sub-algebraic space U of S, the U-abelian algebraic space A, the A-torsor X on U for the étale topology be as in (8.2) so that $Y \times_S U = Y_U$ is the quotient of X by a basic finite étale U-equivalence relation L. It suffices to show that $U = S$.

Factor $A_t' \to Y_t$ as $A_t' \to X_t \to Y_t$ (4.1). The degree $[X_t : Y_t] = d$, which divides $[A_t' : Y_t]$, is prime to the residue characteristics of S. By (3.4), there is a maximal open and closed sub-algebraic space U' of U such that $X_{U'}$ is of constant degree d over $Y_{U'}$. By (8.1), U' is closed in S. So $U' = U = S$, as S is connected.

b) The assertion is immediate by (3.5)+(7.3). And the map which with $P' \in H^1(S, G)$ associates

$$X' = P' \overset{G}{\wedge} X$$

provides the inverse.

□

10. *Irreducibility of non-uniruled degenerate fibers. Almost non-degeneration.*

The irreducibility of non-uniruled degenerate fibers of an abelian fibration is shown in [4], hence :

Proposition 10.1. — *Keep the notations of* (1.1). *Assume that* S *is open in an algebraic space* $\overline{S}$ *and that* Y *is open dense in a separated finitely presented* $\overline{S}$*-algebraic space* $\overline{Y}$ *with structural morphism* $\overline{f}$. *Let* $\overline{s}$ *be a geometric point of* $\overline{S}$ *with values in an algebraically closed field.*

Then $\overline{f}^{-1}(\overline{s})$ *is irreducible if it is non-empty and does not have uniruled irreducible components and if one of the following two conditions holds* :

a) $\overline{f}$ *is flat at all maximal points of* $\overline{f}^{-1}(\overline{s})$.

b) $\overline{S}$ *is locally noetherian and* $\overline{f}$ *is universally open at all maximal points of* $\overline{f}^{-1}(\overline{s})$.

Proof. By standard arguments one may assume that $\overline{S}$ is the spectra of a complete discrete valuation ring with $\overline{S} - S = \{\overline{s}\}$, that $\overline{f}$ is flat and that $\overline{f}^{-1}(\overline{s})$ has no imbedded components. Let X be as in (1.1) and $\overline{X}$ the normalization of $\overline{Y}$ in X. Then $\overline{X}_{\overline{s}}$ is non-empty and does not have uniruled irreducible components and hence by [4] 4.1 is irreducible. So $\overline{f}^{-1}(\overline{s})$ is irreducible.

□

Theorem 10.2. — *Let* S *be the spectra of a discrete valuation ring and* $\overline{t}$ *(resp.* $\overline{s}$*) a geometric generic (resp. closed) point of* S *with values in an algebraically closed field. Let* Y *be an* S*-algebraic space with structural morphism* f. *Assume that* f *is separated of finite type, that* Y *is normal integral and at each of its geometric codimension* ≥ 2 *points either regular or pure geometrically para-factorial of equal characteristic, that* $f^{-1}(\overline{t})$ *is the quotient of a* $\overline{t}$*-abelian variety* $A'_{\overline{t}}$ *by a finite étale* $\overline{t}$*-equivalence relation with degree* $[A'_{\overline{t}} : f^{-1}(\overline{t})]$ *prime to the characteristic of* $k(\overline{s})$ *and that* $f^{-1}(\overline{s})$ *is non-empty, proper, of total multiplicity prime to the characteristic of* $k(\overline{s})$ *and does not have uniruled irreducible components.*

Then f *factors canonically as* $Y \to E \to S$ *with* $Y \to E$ *proper and smooth, where* E *is a finite flat* S*-algebraic stack, regular, tame over* S *and satisfies* $E \times_S \overline{t} = \overline{t}$.

Proof. Notice that f is faithfully flat and, being separated of finite type with geometrically irreducible (10.1) and proper fibers, that f is also

proper (EGA IV 15.7.10). Let the total multiplicity of $f^{-1}(\overline{s})$, that is by definition, the greatest common divisor of the lengths of the local rings of $f^{-1}(\overline{s})$ at its maximal points, be δ, which by hypothesis is prime to the characteristic of $k(\overline{s})$. Thus, the 1-codimensional cycle on Y with rational coefficients, $\Delta = f^*\mathrm{Cyc}_S(\pi)/\delta$, where π is a uniformizer of S, has integral coefficients and is a prime cycle and is locally principal, for Y, being normal and geometrically para-factorial at all its geometric codimension ≥ 2 points, has geometrically factorial local rings (EGA IV 21.13.11). With Δ one associates a canonical μ_δ-torsor on Y for the étale topology, $Y' \to Y$. Let $S' = \mathrm{Spec}\ \Gamma(Y', \mathcal{O}_{Y'})$. There is by quotient by μ_δ an S-morphism

$$Y = [Y'/\mu_\delta] \to [S'/\mu_\delta] = E.$$

It suffices evidently to show that $Y' \to S'$ is smooth, for then

$$Y \to E \to S$$

is the desired factorization of f. Replacing f by $Y' \to S'$, we assume from now on that $\delta = 1$, namely, that $f^{-1}(\overline{s})$ is integral. And, replacing f by $f \times_S S_{(\overline{s})}$, where $S_{(\overline{s})}$ is the strict henselization of S at $\overline{s}$, we assume that S is strictly henselian.

Let t (resp. s) be the generic (resp. closed) point of S. Choose by (1.1) a triple (A_t, X_t, L_t) so that, for a t-abelian variety A_t, $Y_t = f^{-1}(t)$ is the quotient of an A_t-torsor X_t by a basic finite étale t-equivalence relation L_t. Let $G_t = f_{t*}\underline{\mathrm{Aut}}_{Y_t}(X_t)$. The degree $[X_t : f^{-1}(t)] = [G_t : t]$, which divides $[A'_{\overline{t}} : f^{-1}(\overline{t})]$ (4.1), is prime to the characteristic of $k(s)$.

Let X be the normalization of Y in X_t. Then $X_{\overline{s}}$ is non-empty and does not have uniruled irreducible components and hence is irreducible (10.1). Let $p : X \to Y$ be the projection and x (resp. $y = p(x)$) the generic point of X_s (resp. Y_s). Note that there is an open neighborhood of x (resp. y) in X (resp. Y) which is a scheme ([2] 3.3.2).

— *Reduction to the case where G_t is constant and cyclic* :

One has that $[X_t : Y_t] = [\mathcal{O}_{X,x} : \mathcal{O}_{Y,y}] = e[k(x) : k(y)]$ is prime to the characteristic of $k(s)$, where e is the ramification index of $\mathcal{O}_{X,x}$ over $\mathcal{O}_{Y,y}$. One deduces that there is an open sub-scheme V of Y containing y such that V is S-smooth, that $(U_s)_{\mathrm{red}}$ is finite étale surjective over V_s of rank $[k(x) : k(y)]$, where $U = p^{-1}(V)$, and that the ideal of U defining the closed sub-scheme $(U_s)_{\mathrm{red}}$ is generated by one section $h \in \Gamma(U, \mathcal{O}_U)$. In particular, U is regular with h_u being part of a regular system of parameters at each point u of U_s. Now S being strictly henselian, choose a point $u \in U_s(s)$, let $v = p(u)$, let $n = \dim_u(U_s)$ and choose $h_1, \cdots, h_n \in \mathcal{O}_{U,u}$ so that $\{h_1 \bmod h, \cdots, h_n \bmod h\}$ is the

image of a regular system of parameters of V_s at v. Then $h, h_1, \cdots, h_n$ form a regular system of parameters of U at u. Let

$$R = \mathrm{Spec}(\mathcal{O}_{U,u}/(h_1, \cdots, h_n)),$$

which is regular local of dimension 1 and finite flat tame along s of rank e over S. Let r be the generic point of R. The closed image of $p(r)$ in Y is an S-section lying in V and one obtains the following commutative diagram of S-schemes :

$$\begin{array}{ccc} R & \longrightarrow & U = p^{-1}(V) \\ \downarrow & & \downarrow p \\ S & \longrightarrow & V \end{array}$$

The G_t-torsor structure on $p^{-1}(p(r))$, where one identifies the t-rational point $p(r)$ with t, induces an epimorphism

$$G_t \times_t r \to p^{-1}(p(r)),\ (g, \lambda) \mapsto g.\lambda$$

and an isomorphism

$$G_t/\underline{\mathrm{Norm}}_{G_t}(r) \xrightarrow{\sim} \pi_o(p^{-1}(p(r))).$$

Let $Z_t = X_t/N$ be the quotient of X_t by $N = \underline{\mathrm{Norm}}_{G_t}(r)$ and let Z be the normalization of Y in Z_t. The fiber Z_s is irreducible with generic point z being the image of x. Observe that, on writing w for the image of u in Z, Z is by construction étale over Y at w and *a priori* at z. So $Z \to Y$ is étale, as Y by hypothesis is pure at all its geometric codimension ≥ 2 points. Observe next that N is constant and cyclic. Replacing Y by Z and G_t by N, one may assume that $G_t \simeq \mathbf{Z}/e\mathbf{Z}$.

— *Assume $G_t = \mathbf{Z}/e\mathbf{Z}$. Reduction to the case where $p : X \to Y$ is étale* :

Identify $\mathbf{Z}/e\mathbf{Z}$ with μ_e. The μ_e-torsor $X_t \to Y_t$ corresponds to a pair (J_t, α_t) which consists of an invertible module J_t on Y_t and of an isomorphism $\alpha_t : \mathcal{O}_{Y_t} \xrightarrow{\sim} J_t^{\otimes e}$. There is an invertible $\mathcal{O}_Y$-module J extending J_t, since Y has geometrically factorial local rings. Since furthermore f has geometrically integral fibers, $J^{\otimes e}$ is isomorphic to $\mathcal{O}_Y$, say by $\beta : \mathcal{O}_Y \xrightarrow{\sim} J^{\otimes e}$. The difference in $H^1(Y_t, \mu_e)$ of the classes of (J_t, α_t) and (J_t, β_t), that is, the class of $(\mathcal{O}_{Y_t}, \beta_t^{-1}\alpha_t)$, is contained in the image of the map

$$f^* : H^1(t, \mu_e) \to H^1(Y_t, \mu_e),$$

for one has $\Gamma(Y_t, \mathcal{O}_{Y_t}) = k(t)$. By (1.1) *b*) replacing X_t by the μ_e-torsor $X_t' \to Y_t$ defined by (J_t, β_t), one may assume that $p : X \to Y$ is étale. It suffices to show that X is smooth over S. This follows from [4] 4.3. □

Lemma 10.3. — *Keep the notations of* (1.1). *Assume that S is a noetherian local scheme with closed point s such that one of the following two conditions holds* :

a) S is regular of dimension > 0.

b) S is pure geometrically para-factorial along s of equal characteristic.

Let $U = S - \{s\}$. Then each U-section of $f|U$ extends uniquely to an S-section of f.

Proof. In case a) one applies [4] 2.1 as the geometric fibers of f do not contain rational curves. Assume b). One may assume S strictly local. Choose (A, X, L) as in (1.1), let $G = f_*\underline{\mathrm{Aut}}_Y(X)$, $p : X \to Y$ the projection and $y : U \to Y$ a section of $f|U$. The finite étale S-group G is constant and the $G|U$-torsor $p^{-1}(y)$ is trivial, for S is strictly local and pure along s. By *loc.cit.* 5.2+5.3 each U-section of $p^{-1}(y)$ extends uniquely to an S-section of X, hence the claim.

□

Lemma 10.4. — *Let S be a noetherian normal local scheme of equal characteristic zero pure geometrically para-factorial along its closed point s. Let $U = S - \{s\}$. Let E be the fiber category on the category of S-algebraic spaces whose fiber over each S-algebraic space S' is the full sub-category of the category of S'-algebraic spaces consisting of the S'-algebraic spaces Y' which, for an S'-abelian algebraic space A', an A'-torsor X' on S' for the étale topology of finite order and a basic finite étale S'-equivalence relation L' on X', is representable as the quotient X'/L'.*

Then the restriction functor $E(S) \to E(U)$ is an equivalence of categories.

Proof. This restriction functor is fully faithful by (10.3), since every S-smooth algebraic space is pure geometrically para-factorial along its closed S-fiber.

This functor is essentially surjective. For, U-abelian algebraic spaces extend to S-abelian algebraic spaces ([4] 5.1). And, as S is pure along s, if n is an integer ≥ 1, A an S-abelian algebraic space and ${}_nA = \mathrm{Ker}(n.\mathrm{Id}_A)$, ${}_nA|U$-torsors on U for the étale topology extend to ${}_nA$-torsors on S for the étale topology. Thus, every object of $E(U)$ is a quotient $(X|U)/(G|U)$ where, for an S-abelian algebraic space A, X is an A-torsor on S for the étale topology of finite order and G is a finite étale S-group such that $G|U$ acts and defines a basic finite étale U-equivalence relation on $X|U$. Each such action $G|U \times_U X|U \to X|U$

extends by [4] 5.2+5.3 to a unique S-morphism $\mu : G \times_S X \to X$, since $G \times_S X$ is geometrically para-factorial along its closed S-fiber. Clearly, μ represents a G-action and defines a basic finite étale S-equivalence relation on X. And X/G is the desired extension of $(X|U)/(G|U)$.

□

Theorem 10.5. — *Let S be an integral scheme with generic point t. Let F be the fiber category on the category of S-algebraic spaces whose fiber over each S-algebraic space S' is the full sub-category of the category of S'-algebraic spaces consisting of the S'-algebraic spaces Y' which, for an S'-abelian algebraic space A', an A'-torsor X' on S' for the étale topology and a basic finite étale S'-equivalence relation L' on X', is representable as the quotient X'/L'. Let Y be an S-algebraic space with structural morphism f. Assume that Y is locally noetherian normal integral of residue characteristics zero and at all its geometric codimension ≥ 2 points pure and geometrically para-factorial. Assume furthermore that $f^{-1}(t)$ is an object of $F(t)$ and that, for each geometric codimension 1 point $\overline{y}$ of Y, $f \times_S S_{(\overline{s})}$ is separated of finite type and flat at $\overline{y}$ and the geometric fiber $f^{-1}(\overline{s})$ is proper and does not have uniruled irreducible components, where $S_{(\overline{s})}$ denotes the strict localization of S at $\overline{s} = f(\overline{y})$.*

Then up to unique isomorphisms there exists a unique groupoid in the category of S-algebraic spaces whose nerve $(Y_\cdot, d_\cdot, s_\cdot)$ satisfies the following conditions :

a) $Y = Y_o$.

b) The Y-algebraic space with structural morphism $d_1 : Y_1 \to Y_o$ is an object of $F(Y)$.

c) Over t, $Y_{\cdot t} = \mathrm{cosq}_o(f^{-1}(t)/t)$.

Proof. Notice that for each geometric codimension 1 point $\overline{y}$ of Y the localization of S at the image s of $\overline{y}$ is noetherian regular of dimension ≤ 1, since f is by hypothesis flat at $\overline{y}$. If S is local of dimension 1 with closed point s, then with the notations of (10.2) the asserted S-groupoid is $\mathrm{cosq}_o(Y/E)$ which, as Y is regular (10.2), is unique up to unique isomorphisms (10.3). By a "passage à la limite" and by gluing, one obtains in the general case an S-groupoid $U_\cdot$ satisfying the following conditions :

a) U_o is open in Y such that $\mathrm{codim}(Y - U_o, Y) \geq 2$.

b) The U_o-algebraic space with structural morphism $d_1 : U_1 \to U_o$ is an object of $F(U_o)$.

c) Over t, $U_{\cdot t} = \mathrm{cosq}_o(f^{-1}(t)/t)$.

As $d_1 : U_1 \to U_o$ has the canonical section $s_o : U_o \to U_1$, one may by the proof of (8.2) write d_1 as a quotient by a basic finite étale U_o-equivalence relation on a U_o-abelian algebraic space with s_o being the image of the zero section. Hence, by (10.4) there is a cartesian diagram of S-algebraic spaces

$$\begin{array}{ccc} U_1 & \longrightarrow & Y_1 \\ {\scriptstyle d_1}\downarrow & & \downarrow{\scriptstyle d_1} \\ U_o & \longrightarrow & Y_o \end{array}$$

where $Y_o = Y$ whose vertical arrow on the right is an object of $F(Y_o)$. By (10.3) this diagram is unique up to unique isomorphisms and there is a unique extension of $s_o : U_o \to U_1$ to a section $s_o : Y_o \to Y_1$ of $d_1 : Y_1 \to Y_o$. By again (10.3) and arguing as in [4] 5.7 one finds a unique S-groupoid $Y_.$ with $d_1 : Y_1 \to Y_o$ being a face.

□

Similarly as [4] 3.1, we call every groupoid $Y_.$ satisfying the conditions (10.5) a)+b) an *almost non-degenerate fibration structure* on $f : Y \to S$ with $[Y_.]$ being called the *ramification S-stack*. Each such fibration has again a tautological factorization

$$f : Y \to [Y_.] \to S.$$

We say that this structure is *non-degenerate* if the groupoid $Y_.$ is simply connected with $\mathrm{Coker}(d_o, d_1) = S$, that is, if $f : Y \to S$ is an object of $F(S)$ of (10.5). Note that $Y \to [Y_.]$ is then *non-degenerate* in this sense as in *loc.cit.*

Proposition 10.6. — *Keep the notations of* (10.5). *Consider the following conditions* :

1) *f is proper, S is excellent regular.*

2) *f is proper, S is locally noetherian normal and at each of its points satisfies the condition* (W) (EGA IV 21.12.8).

Then, if 1) *(resp.* 2)*) holds, S is the cokernel of (d_o, d_1) in the full sub-category of the category of S-algebraic spaces consisting of the S-algebraic spaces (resp. S-schemes) which are S-separated and locally of finite type over S.*

Proof. One argues as in [4] 5.11.

□

References

1. D. Mumford. Geometric Invariant Theory. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, 1965.

2. M. Raynaud. Spécialisation du foncteur de Picard. *Publications Mathématiques de l'IHÉS*, 38, 1970.
3. M. Raynaud. Faisceaux amples sur les schémas en groupes et les espaces homogènes. *Lecture Notes in Mathematics*, 119, 1971.
4. Y. Zong. Almost non-degenerate abelian fibrations. arxiv.org/abs/1406.5956.

Department of Mathematics, University of Toronto
E-mail address: zongying@math.utoronto.ca